

You Are Cordially Invited to Our Scientific Symposium at Euroanaesthesia 2019 in Vienna

Protecting Vital Organs in High Risk Patients - The Present & Future Role of Noninvasive Smart Technology in Advancing Patient Safety



Location: Strauss 1 Room, Messe Wien Congress Center
Vienna, Austria

Date and Time: Sunday June 2nd • 12:15 - 13:45
Lunch will be provided

Chairperson: Basil Matta, MD, BA, BAO, BCh, MB, FRCA, FFICM

[Please click here to register.](#)



Presenters



Caring for High Risk Patients - New Insights, New Challenges in Optimizing Vital Organ Monitoring

Basil Matta, MD, BA, BAO, BCh, MB, FRCA, FFICM

Divisional Director, MSK. Digestive Diseases, Major Trauma and Perioperative Care Medicine
Clinical Lead - Cambridge University Hospitals Trust
Consultant, Anaesthesia and Critical Care
Associate Lecturer, University of Cambridge
Cambridge, United Kingdom



Hypoxaemia or Hyperoxia - Which Is Worse? - New Horizons

Sigismund Lasocki, MD, PhD

Professor of Anesthesiology and Intensive Care
Chief Department of Anesthesiology and Intensive Care
Chairman - Pôle ASUR (Anesthésie SAMU Urgences Réanimation), CHU Angers
Angers, France



Optimizing Oxygen Delivery: Do We Need to Redefine Our Haemodynamic Goals?

Azriel Perel, MD

Professor of Anesthesiology and Intensive Care
Department of Anesthesiology and Intensive Care
Sheba Medical Center, Tel Aviv University
Tel Aviv, Israel



Cerebral Oxygenation: New Insights into the Mechanism of Delirium in Cardiac Surgery - Reasons for a Change!

André Denault, MD, PhD, ABIM-CCM, FRCPC, FASE, FCCS

Professor of Anesthesiology and Intensive Care
Department of Anesthesiology, Critical Care Program
Montreal Heart Institute, and Centre Hospitalier de l'Université de Montréal
Université de Montréal
Montreal, Quebec, Canada



Post-operative Cognitive Dysfunction (POCD) & Delirium: Preventing Damage and Improving Outcome - The Role of Bilateral EEG

Massimo Lamperti, MD, MBA

Clinical Professor of Anesthesiology
Acting Institute Chair
Anesthesiology Institute-Cleveland Clinic Abu Dhabi
Cleveland Clinic Lerner College of Medicine of Case Western Reserve University
Abu Dhabi, United Arab Emirates



Interactive Session, please ask any questions to our Faculty now! For more information, please stop by Masimo,
Stand #B0401. [Please click here to register and ask your questions.](#)

Masimo may be required to report the value of the meal, or other transfers of value, as required by applicable federal, state or local laws and regulations (e.g., U.S. Physician Payment Sunshine Act, France Sunshine Act).

Review Article

CPD available at <http://www.learnataagbi.org>

Postoperative sore throat: a systematic review

K. El-Boghdadly,¹ C. R. Bailey² and M. D. Wiles³

1 Anesthesia Fellow, Department of Anesthesia, Toronto Western Hospital, Toronto, Ontario, Canada

2 Consultant, Department of Anaesthetics, Evelina London Children's Hospital, Guys and St. Thomas' NHS Foundation Trust, London, UK

3 Consultant, Department of Anaesthetics, Sheffield Teaching Hospital NHS Foundation Trust, Sheffield, UK

Summary

Postoperative sore throat has a reported incidence of up to 62% following general anaesthesia. In adults undergoing tracheal intubation, female sex, younger age, pre-existing lung disease, prolonged duration of anaesthesia and the presence of a blood-stained tracheal tube on extubation are associated with the greatest risk. Tracheal intubation without neuromuscular blockade, use of double-lumen tubes, as well as high tracheal tube cuff pressures may also increase the risk of postoperative sore throat. The expertise of the anaesthetist performing tracheal intubation appears to have no influence on the incidence in adults, although it may in children. In adults, the i-gel™ supraglottic airway device results in a lower incidence of postoperative sore throat. Cuffed supraglottic airway devices should be inflated sufficiently to obtain an adequate seal and intracuff pressure should be monitored. Children with respiratory tract disease are at increased risk. The use of supraglottic airway devices, oral, rather than nasal, tracheal intubation and cuffed, rather than uncuffed, tracheal tubes have benefit in reducing the incidence of postoperative sore throat in children. Limiting both tracheal tube and supraglottic airway device cuff pressure may also reduce the incidence.

Correspondence to: K. El-Boghdadly

Email: elboghdadly@gmail.com

Accepted: 2 February 2016

Keywords: laryngeal masks; postoperative complications; sore throat; supraglottic airway devices; tracheal tubes

Introduction

Postoperative sore throat is a common occurrence following general anaesthesia and, although clinicians often regard it as a relatively minor complication, patients perceive avoidance as being of great importance. Instrumentation of the airway is an inherent risk factor for the development of this common complication, yet a number of techniques can reduce the incidence. We performed a systematic review in order to assess the risk factors that increase the likelihood of postoperative sore throat in adults and children and the efficacy of interventions designed to reduce its incidence.

Methods

We independently and systematically searched the literature using MEDLINE and EMBASE bibliographic databases, the Cochrane Central Register of Controlled Trials (CENTRAL) and manually using the following search terms in either the title or abstract: endobronchial intubation; endotracheal intubation; nasotracheal intubation; intubation, intra-tracheal; supraglottic airway(s) device(s); LMA laryngeal mask Classic; Flexible LMA laryngeal mask; LMA laryngeal mask Unique; Portex SoftSeal; Ambu AuraOnce; Cobra perilaryngeal; cobraPLA; ProSeal LMA laryngeal mask; LMA laryngeal mask Supreme; LMA laryngeal mask Guardian;

g-LMA laryngeal mask; SLIPA; streamlined liner of the pharyngeal airway; i-gel; supraglottic gel device(s); sore throat and pharyngitis. Individual searches were, subsequently combined using AND/OR options. The following filters were then applied: publication date 01/01/2005 to 31/12/2015; English language; human subjects; adult subjects (age ≥ 18 years) and clinical trial. For the section on paediatrics, the same search was performed with the exclusion of adult patients (age < 18 years). Any trials that did not have postoperative sore throat as a primary outcome measure were not included (Fig. 1). These were then qualitatively supplemented by a manual search to include studies that were deemed relevant.

Postoperative sore throat following tracheal intubation in adults

Risk factors

Tracheal intubation is associated with a greater risk of postoperative sore throat than when either a supraglottic airway device or a facemask is used [1] and several risk factors have been identified. A prospective study of 809 patients found a 40% incidence [2]. Subsequent

logistic regression analysis demonstrated that female sex [odds ratio (OR) 1.66], pre-existing lung disease (OR 3.12), duration of anaesthesia (OR 1.27) and the presence of a blood-stained tracheal tube on extubation (OR 4.81) were all associated with the greatest risk of postoperative sore throat. In addition, age was inversely related to the risk [3]. Women may be more likely to report any postoperative complication, thereby potentially introducing reporting bias, which could account for the higher incidence seen in women [4]. A more likely cause, however, are the varying sizes of tracheal tubes used in trials and clinical practice. In a study by Biro et al., [2] 7.5-mm internal diameter (ID) tracheal tubes were used, and it is possible that they were too large for certain patients. This theory is supported by a study in which 89% of female patients underwent tracheal intubation with a 6.0-mm ID tracheal tube. The rate of postoperative sore throat was similar to that seen in the male patients, of whom 97% had an 8.0-mm ID tracheal tube (27% vs 38%, respectively, $p = 0.20$) [5]. A similar reduction in the incidence was seen in a randomised controlled trial that included 100 female patients when a 6.0-mm ID rather than a 7.0-mm ID tracheal tube was used (27.1% vs 51.1%, respectively) [6]. In addition, a subsequent meta-analysis of trials involving 509 female patients suggested that the use of 6.0 mm rather than 7.0-mm ID tracheal tubes reduced the incidence of postoperative sore throat both in the recovery unit and at 24 h postoperatively, with risk reductions of 0.56 and 0.69, respectively [7]. It may appear somewhat surprising that a 1-mm reduction in internal diameter should result in such a marked decrease in the rate of sore throat, since it is only the tracheal tube cuff that is in contact with the trachea. However, what patients describe as 'sore throat' encompasses a wide range of conditions including pharyngitis, laryngitis, tracheitis, cough, hoarseness or dysphagia. Smaller tracheal tubes subjectively provide a better view of the passage of the tube through the larynx, which may reduce the trauma associated with laryngoscopy and tube insertion [8].

The GlideScope[®] (Verathon Inc., Bothell, WA, USA) [9] and the Airway Scope (Pentax, Tokyo, Japan) [10] have both been shown to be associated with lower rates of postoperative sore throat when compared with direct laryngoscopy using Macintosh




Figure 1 Study selection flow diagram.

blades. However, in both these studies, relatively large diameter tracheal tubes were used in female patients (7.0-mm ID and 7.5-mm ID) and tracheal cuff pressures were not accurately measured or monitored. The expertise of the anaesthetist performing tracheal intubation appears to have no effect as, in a retrospective review of 21,606 patients [3], consultants and trainees had a similar incidence of throat complications (32.6% vs. 32.9%, respectively).

Undertaking tracheal intubation without neuromuscular blockade increases the incidence of postoperative sore throat [11]. It has been suggested that the use of suxamethonium may increase sore throat, possibly due to fasciculation and subsequent myalgia of striated pharyngeal muscle fibres. A recent study, however, found a similar incidence and severity following the use of suxamethonium and rocuronium [12].

Double-lumen endobronchial tubes are associated with a greater risk of sore throat (OR 2.55) [13], although those made from silicon may have lower rates than polyvinylchloride [14]. In addition, insertion with the tracheal lumen facing anteriorly during passage through the vocal cords reduces rates by 50% [15]. A recent systematic review and meta-analysis demonstrated that bronchial blockers are associated with a lower incidence compared with double lumen tubes (OR 0.39) [13]; all types of bronchial blocker appear to be similarly efficacious in this regard [16]. The type of single-lumen tracheal tube has no effect [17].

Prevention

Lidocaine: A meta-analysis by the Cochrane Collaboration [18] reviewed 19 studies involving 1,940 patients that used lidocaine in several different ways and in a variety of concentrations: within the tracheal tube cuff (2–10%); applied as a gel on the exterior of the tracheal tube (4%); intravenously (1–1.5 mg.kg⁻¹) or as an aerosol spray (4–10%). Overall, topical and systemic lidocaine therapy appeared to reduce the risk of postoperative sore throat [risk ratio (RR) 0.64, 95% confidence interval (CI) 0.48–0.85]; however, when only the studies rated as being of high quality were analysed, this benefit was no longer observed (RR 0.71, 95% CI 0.47–1.09). Due to variations in study methodology and the differing doses and routes of administration, we cannot recommend lidocaine for

the prevention of postoperative sore throat, and some additives in aerolised lidocaine may actually cause it [18]. Since the publication of the meta-analysis, three further studies have been published, involving a total of 370 patients, which have suggested that lidocaine is of benefit when directly applied to the glottis [19, 20] or administered within the tracheal tube cuff [21]. However, these studies are of low quality and the conclusions of the Cochrane group remain valid.

Steroids: Dexamethasone is the most popular steroid studied in this regard. A meta-analysis published in 2014 that included seven randomised controlled trials suggested that an intravenous dose greater than 0.1 mg.kg⁻¹ reduced the incidence and severity of postoperative sore throat at 24 h (RR 0.68 and standardised mean difference -1.15) [22]. The smallest tracheal tube used in these studies was 7.0-mm ID, one study used only double lumen tubes and tracheal cuff pressures were controlled in only one study. The application of triamcinolone paste (0.1%) to the tracheal tube and cuff was associated with a reduction in the incidence and severity of sore throat at 24 h compared with chlorhexidine gel [23]. Similarly, betamethasone gel (0.05%) reduced the incidence compared with 2% lidocaine gel [24]. Pre-operative inhaled fluticasone [25] and budesonide [26] have also been shown to reduce the incidence and severity of postoperative sore throat.

Non-steroidal anti-inflammatory drugs (NSAIDs): The NSAID most commonly investigated is benzydamine hydrochloride, a topical preparation that also has antibacterial properties. A meta-analysis of benzydamine identified five randomised controlled trials that included 824 patients, one of which used supraglottic airway devices [27]. For the studies investigating tracheal tubes, benzydamine was sprayed directly onto the tracheal tube and/or into the oropharynx in three studies while the remaining study used a benzydamine gargle. The incidence of postoperative sore throat compared with control groups was reduced at all measured time points up to 24 h (RR at 1 h and 24 h were 0.42 and 0.32, respectively). A more recent study examined the effect of applying benzydamine to the oropharynx and

tracheal tube cuff of patients requiring double lumen tracheal tubes [28]. The incidence and severity of postoperative sore throat was reduced at 1 h, 6 h and 24 h; however, the reduction in severity of symptoms was modest, with a mean improvement in VAS scores of less than 13 mm in a scale of 0–100 mm. A pre-operative 350 mg aspirin gargle reduced postoperative sore throat, but only for 2 h [29]. Intravenous diclofenac had no effect in patients undergoing laparoscopic surgery [30], although topical diclofenac reduced the incidence of sore throat following caesarean section [31].

Tracheal tube cuff pressure: The monitoring and limitation of tracheal tube cuff pressure as a method to reduce postoperative sore throat has been extensively investigated because excessive cuff pressure can damage tracheal mucosa by direct trauma and reduction in blood flow. A prospective, randomised controlled trial compared tracheal tube cuff inflation using a manometer (pressure 15–25 mmHg) with manual palpation of the pilot balloon in 509 patients [32]. The cuff pressure, however, was not subsequently monitored during the remainder of surgery. The manometer group patients had a significant reduction in the incidence of sore throat at 24 h compared with control (34% vs 44%, respectively). Subsequent studies in patients having maxillofacial [33] and thyroid surgery [34] have confirmed that control of tracheal tube cuff pressure reduces the severity of symptoms for 2–6 h and the incidence of sore throat for up to 24 h.

Liquorice: Liquorice is derived from the root of *Glycyrrhiza glabra* and has been utilised in traditional and modern medicine because some of its ingredients are reported to have anti-inflammatory and antitussive properties. Agarwal et al. demonstrated a reduction in the incidence of sore throat for up to 4 h postoperatively following a gargle of 0.5 g liquorice in 30 ml water 5 min before induction of anaesthesia [35]. However, at 24 h postoperatively the incidence of pain at rest was the same as in the placebo group. In a larger cohort ($n = 236$) who underwent tracheal intubation with double lumen endobronchial tubes, the incidence of sore throat was reduced by around 50% in comparison with controls (21% vs. 45%,

respectively; RR 0.48) [36]. Similar benefits were found in smokers who sucked liquorice lozenges 30 min before surgery [37].

N-methyl-D-aspartate (NMDA) receptor antagonists: The antinociceptive and anti-inflammatory effects of the NMDA receptor antagonists, magnesium and ketamine, have been investigated in attenuating the symptoms of postoperative sore throat. The sucking of a lozenge containing 610 mg magnesium citrate 30 min before surgery significantly reduced the incidence at 2 h and 4 h postoperatively when compared with a placebo lozenge (14% vs 40%, respectively at 4 h) [38]. By 24 h, however, the incidence of sore throat was similar in both groups. A magnesium gargle (20 mg.kg^{-1}) administered 15 min before appendectomy reduced the incidence and severity of postoperative sore throat at 2 h, 4 h and 24 h compared with a ketamine gargle (0.5 mg.kg^{-1}) [39], although a pre-operative ketamine gargle (40 mg) was effective for 24 h in another study [40]. The mechanism of action of ketamine is likely due to a local effect, because an intravenous 0.5 mg.kg^{-1} bolus and subsequent infusion had no effect on the incidence or severity of sore throat 24 h following cholecystectomy [41].

Others: Several other novel methods have been examined, including lozenges containing amyl-m-cresol (the active ingredient in Strepsils[®]), azulene and dexpanthenol [42–45]. Although the results of these trials all reported reductions in the incidence of postoperative sore throat, larger studies are needed to confirm the efficacy and safety of these agents. Premedication with oral clonidine $150 \mu\text{g}$ is ineffective and may exacerbate symptoms, presumably secondary to an antisialogogue effect [46]. Gabapentin 600 mg, 1 h before surgery reduced the incidence of sore throat at rest but not on swallowing in patients who had undergone thyroid surgery [47].

Postoperative sore throat following supraglottic airway device (SAD) use in adults

Supraglottic airway devices are used in approximately 56% of all general anaesthetics administered in the UK, of which 80% are first generation devices [48]. First-generation SADs lack design features to reduce

the risk of gastric aspiration while second-generation SADs offer improved safety against gastric aspiration and regurgitation, often have integral bite blocks and provide higher oropharyngeal leak pressures. Although the incidence of postoperative sore throat is lower compared with tracheal tubes, it still remains significant at up to 49% [49] and is affected by the choice of SAD, insertion technique and post-insertion management. Commonly performed trials compare the clinical characteristics of two or more devices; however, the primary outcome measure in the vast majority of studies is not postoperative sore throat. Although our search strategy identified two studies [50, 51] comparing the effect of SADs on postoperative sore throat as the primary outcome measure, for the purposes of comparison, we undertook a manual search and selection of studies we felt were relevant comparators. In these, the incidence of postoperative sore throat for each device varies markedly between trials; for example, the LMA[®] laryngeal mask¹ Classic (Teleflex Medical, Morrisville, NC, USA) has a reported incidence between 2.6% [52] and 42% [53]. This disparity is probably attributable to differences in quantifying postoperative sore throat and/or different study methodology.

Choice of SAD

When comparing the LMA laryngeal mask Classic[™] with the single use Portex[®] Soft Seal[®] (Smiths Medical, St. Paul, MN, USA), Tan et al. [54] found no difference in postoperative sore throat incidence, although this was in contrast to a number of older studies [55–57]. When comparing the Soft Seal[®] and the LMA laryngeal mask Unique[™] (Teleflex Medical), one study found a higher incidence in the Soft Seal group [58], while another found no difference [59]. The Cobra perilaryngeal airway (CobraPLA[®], Pulmo-dyne, Indianapolis, IN, USA) also has variable results [60–62] and it is likely that there is little difference in postoperative sore throat incidence when comparing first-generation SADs.

Second-generation SADs have a more compelling evidence base. Numerous studies have demonstrated a

reduction in postoperative sore throat when using the i-gel[™] (Intersurgical, Wokingham, UK) [50, 63–65], although few demonstrated equivalent outcomes to other SADs [66]. A meta-analysis found no difference in the incidence between the ProSeal LMA laryngeal mask[™] (Teleflex Medical, NC, USA) and the LMA laryngeal mask Supreme[™] (Teleflex Medical) [67]. The ProSeal LMA laryngeal mask has a similar sore throat profile to that of the LMA laryngeal mask Guardian[™] (Teleflex Medical) [68], and possibly the i-gel [69, 70] but has a variable evidence base when compared with the streamlined liner of the pharynx airway (SLIPA[™], CurveAir Ltd, London, UK) [71, 72]. There was no significant difference with the LMA laryngeal mask Supreme and the SLIPA compared with the LMA laryngeal mask Classic[™] [73, 74]. The incidence of postoperative sore throat with the Ambu[®] AuraOnce[™] (Ambu, Ballerup, Denmark) is also comparable to first generation SADs, including the LMA laryngeal mask Classic, LMA laryngeal mask Unique and the Soft Seal LMA laryngeal mask [75], but higher than the i-gel [76].

The available evidence suggests that there is little difference in the incidence of postoperative sore throat between first- and second-generation SADs, with the exception of the i-gel, possibly due to the absence of an inflatable cuff.

Insertion technique

Multiple modifications in the recommended LMA laryngeal mask Classic insertion technique (placing a fully deflated, lubricated device with the index finger) have been attempted in order to increase success rate and reduce the incidence of laryngopharyngeal complications. Supraglottic airway device insertion with a fully inflated, rather than a fully deflated, cuff was found to reduce the incidence of postoperative sore throat, with less blood staining on removal of the device [77]. An alternative strategy involves inserting a deflated SAD without using an index finger but, although a more appealing technique, this has a similar insertion success rate to the manufacturers' recommended method and the same incidence of postoperative sore throat [78]. Interestingly, inflating the SAD cuff *after* securing the device has been shown to reduce the incidence when compared with the traditional method of securing the device after inflating the

¹ LMA is a registered trade mark of The Laryngeal Mask Company Ltd, an affiliate of Teleflex Incorporated.

cuff [79]. In one study, a laryngoscope-guided technique to insert flexible SADs reduced the incidence of postoperative sore throat from 35.2% to 16.7% when compared with the standard insertion technique [80]. The ProSeal LMA laryngeal mask is more difficult to insert than the LMA laryngeal mask Classic [81] and different insertion techniques have been developed. With the standard approach, there is a sore throat incidence of up to 33% [82], which was reduced to 25% when an introducer was used [83]. The 90° rotational technique is an alternative method, whereby the device is inserted in the midline, following which the shaft is rotated 90° counter-clockwise until resistance is encountered, then the device is rotated clockwise back to the midline. This has a greater insertion success rate, as well as a significantly lower incidence of postoperative sore throat (between 9% and 12%) [81, 82]. The use of a gum elastic bougie to guide ProSeal LMA laryngeal mask insertion has a lower incidence of sore throat when compared with the finger insertion technique [83, 84], and the same holds true when using a stylet [85]. Alternative insertion methods have not demonstrated a reduction [86, 87].

Cuff pressures

Most manufacturers recommend that SAD cuff pressures should not exceed 60 cmH₂O. An increasing body of evidence suggests that high cuff pressures contribute significantly to laryngopharyngeal complications, including postoperative sore throat. The LMA laryngeal mask Classic is the most studied device and monitoring and maintaining a cuff pressure of 60 cmH₂O or lower is associated with a significant reduction in sore throat [88–93]. The effect of cuff pressure is less obvious in other SADs. For example, a study comparing high and low cuff pressures using the ProSeal LMA laryngeal mask found no statistically significant difference [94]. The CobraPLA, which is larger than the LMA laryngeal mask Classic size for size, was studied and low cuff pressures reduced the incidence of moderate sore throat, although mild and severe pain was the same [95]. Using cuff pressures in the LMA laryngeal mask Supreme of 25 cmH₂O resulted in a reduced incidence of postoperative sore throat [96]. These data are from a small study in patients undergoing laparoscopic surgery (n = 98), but suggest that

pressures significantly lower than 60 cmH₂O may be preferable.

Various alternatives to the recommended water, water-based gel or saline lubricants have been investigated. Nasseri et al. [97] compared lubricating the LMA laryngeal mask Classic with either lidocaine 2% or standard lubricant gel and found no difference in the incidence of postoperative sore throat. A betamethasone 0.05% gel was found to be superior to lidocaine 2% gel [98]. Other studies found no benefit with alternative lubricants [99] or humidification techniques [100].

Others

The pre-operative administration of a flurbiprofen lozenge reduced the severity, although not the incidence, of early postoperative sore throat [101]. A pre-operative 2 mg.kg⁻¹ tramadol gargle reduced both the incidence and severity of sore throat for up to 24 h postoperatively in a small study of 50 patients [102], although 100 mg of intravenous hydrocortisone had no effect [103]. Diffusion of nitrous oxide into the cuff led to an increase in cuff pressure of up to 250% [104] and resulted in an increased incidence of postoperative sore throat [91]. Neuromuscular blockade before the insertion of a ProSeal LMA laryngeal mask doubled the incidence of sore throat [105]. The use of propofol is associated with a better laryngopharyngeal morbidity profile compared with sodium thiopental when using a SAD [106] and has a more favourable outcome when used for anaesthesia maintenance compared with sevoflurane [107].

Postoperative sore throat in children

The Royal College of Anaesthetists (RCOA) state that 'For a child in good health having minor surgery one child in ten experiences a headache or a sore throat' [108]. At the Evelina London Children's Hospital, a postoperative sore throat incidence of between 1% and 10% is quoted [109]. The incidence appears to be lower in children compared with adults, but it is difficult for a small child to specifically express their discomfort, which may manifest as agitation, crying or restlessness and even be misinterpreted as emergence delirium [110] or wound pain [111]. The incidence, as in adults, is much higher if the child or parent/carer is specifically questioned [112]. The severity is traditionally categorised on a four-point scale as described by Stout et al. [113] but this scale is

not very useful in children. It is usually at its worst 4 h postoperatively but may persist for up to four days [114].

Pre-operative risk factors

Pre-operative risk factors such as asthma or dry cough make the patient more susceptible to peri-operative adverse respiratory events, including sore throat [115]. Children with an upper respiratory tract infection are significantly less likely to have sore throat and other postoperative respiratory complications when an SAD is used [116], although a study in children with minor upper respiratory tract infections requiring ophthalmic procedures found no difference when compared with the use of a facemask [117].

Prevention

The pre-operative use of an antisialogogue (glycopyrrolate) resulted in postoperative sore throat in 13.3% of children [118]. Local anaesthetic gel applied to an airway device or spraying of the mouth with local anaesthetic may appear attractive but a commercial lidocaine throat spray caused more postoperative problems than saline spray, again, possibly due to irritant additives within the spray itself [119].

Tracheal tubes

When inserting a tracheal tube, postoperative throat complications are, intuitively, more likely [120]. For example, there may be coughing or movement due to inadequate anaesthesia or neuromuscular blockade,

glottic exposure may be difficult, and airway trauma is more common. Operator experience may play a part rather more so than in adults. Supraglottic airway devices caused less sore throat than nasotracheal tubes in children aged 2–7 years having day-case dental surgery [121]. In a meta-analysis of 16 studies, comparing SADs with a tracheal tube, the overall incidence of postoperative sore throat was 9.8% vs 15.3%, respectively [122]. This contrasts with a quantitative meta-analysis of respiratory complications from 19 studies that compared SAD with tracheal intubation [123]. Although the incidence of desaturation, laryngospasm, coughing and breath holding were lower when a SAD was used to secure the airway, the incidence of sore throat (OR 0.87; 95% CI 0.53–1.44) was similar. Nasotracheal is associated with more sore throat than orotracheal intubation. Following general anaesthesia for dental surgery, children whose tracheal intubations were traumatic were more likely to report postoperative sore throat [124].

In a study of 111 children, 37% of those intubated with an uncuffed tracheal tube complained of sore throat compared with 19% with cuffed tracheal tubes [125]. Furthermore, the incidence of sore throat correlated with increased cuff pressure: 4% at cuff pressures of 11–20 cmH₂O; 20% at 21–30 cmH₂O; 68% at 31–40 cmH₂O; and 96% at cuff pressures of > 40 cmH₂O.

Supraglottic airway devices

The paediatric i-gel has the same design as the adult device and has potential to cause less postoperative

Table 1 Potential risk reduction interventions for postoperative sore throat.

Tracheal intubation	SADs	Children
Smaller tube size [5–7]	Use of i-gel [50, 63–65]	SAD rather than tracheal tube [121, 122]
Video laryngoscopy [9, 10]	90° rotational insertion technique, use of introducing stylet for ProSeal LMA laryngeal mask [81–85]	Oral rather than nasotracheal intubation [124]
Limiting cuff pressure [32–34]	Cuff pressure limitation ≤ 60 cmH ₂ O [88–93, 96]	Cuffed rather than uncuffed tubes [125]
Intravenous, topical or inhaled steroids [22–25]	Topical steroids, NSAIDs, tramadol [98, 101, 102]	Limiting tracheal tube cuff pressure [126]
Topical NSAIDs [27–29, 31]	Propofol induction and maintenance [106, 107]	SAD cuff pressure limitation ≤ 60 cmH ₂ O [130–133]
Liquorice, magnesium and ketamine gargle [35–40]		

NSAIDs, non-steroidal anti-inflammatory drugs; SAD, supraglottic airway device.

sore throat because there is no cuff inflation. The i-gel has been compared with the ProSeal LMA laryngeal mask, the LMA laryngeal mask Classic [126], the LMA laryngeal mask Supreme [127] and the PRO-Breathe[®] silicone disposable SAD (Well Lead Medical Co Ltd, Panyu, China) [128] in both spontaneously breathing and patients who received neuromuscular blockers, but these studies were not sufficiently powered to detect a difference in complication rates. However, Smith and Bailey [129] found that there was no difference in the overall incidence of postoperative sore throat between the i-gel and different SADs when data were pooled.

As with adults, most manufacturers recommend cuff inflation with a maximum volume of air or to a maximum pressure of 60 cmH₂O. Inflation to a pressure less than the manufacturers' recommendations still allows adequate ventilation with minimal leakage [130]. Four hundred children receiving an SAD were studied and 45 (11.25%) developed sore throat, of which 56.5% had cuff pressures exceeding 100 cmH₂O [131]. The incidence was zero when cuff pressures were less than 40 cmH₂O. It has been suggested that, particularly for the smaller sizes, SADs should be deflated following insertion, rather than inflated, in order to avoid cuff hyperinflation [132]. The use of nitrous oxide significantly increases cuff pressures and, in one study, the incidence of postoperative sore throat was 45% compared with 5% in the group receiving air [133].

When using flexible SADs, introducers may be used to facilitate insertion and do not appear to increase the incidence of sore throat [134]. However, using flexible laryngeal masks with a polyvinyl chloride surface was associated with a significantly higher risk when compared with a silicone surface. As with cuffed tracheal tubes, when using cuffed SADs, cuff pressures should be monitored in order to minimise pressure-related airway complications.

Conclusion

Postoperative sore throat is largely self-limiting and most interventions only result in a minor reduction in the severity of symptoms (Table 1). The risk following tracheal intubation is most likely to be minimised by using the smallest tracheal tube practically

possible (especially in female patients) and monitoring cuff pressure. Insertion techniques may affect the incidence of postoperative sore throat when using SADs, but the most significant factor is excessive cuff pressure and, therefore, the i-gel has the most favourable profile in adults. Postoperative sore throat in children is a significant complication and the risk should be explained to the child and parent/carer preoperatively. Use of oral tracheal tubes results in less sore throat than nasal tracheal tubes and cuffed ones are less likely to cause postoperative sore throat compared with uncuffed tubes, provided cuff pressures are monitored.

Competing interests

CRB and MDW are Editors of *Anaesthesia* and this article was externally reviewed. No external funding or other competing interests declared.

References

- Higgins PP, Chung F, Mezei G. Postoperative sore throat after ambulatory surgery. *British Journal of Anaesthesia* 2002; **88**: 582–4.
- Biro P, Seifert B, Pasch T. Complaints of sore throat after tracheal intubation: a prospective evaluation. *European Journal of Anaesthesiology* 2005; **22**: 307–11.
- Inoue S, Abe R, Tanaka Y, Kawaguchi M. Tracheal intubation by trainees does not alter the incidence or duration of postoperative sore throat and hoarseness: a teaching hospital-based propensity score analysis. *British Journal of Anaesthesia* 2015; **115**: 463–9.
- Myles PS, Hunt JO, Moloney JT. Postoperative 'minor' complications. *Anaesthesia* 1997; **52**: 300–6.
- Jaensson M, Gupta A, Nilsson U. Gender differences in sore throat and hoarseness following endotracheal tube or laryngeal mask airway: a prospective study. *BMC Anesthesiology* 2014; **14**: 56.
- Jaensson M, Olowsson LL, Nilsson U. Endotracheal tube size and sore throat following surgery: a randomized-controlled study. *Acta Anaesthesiologica Scandinavica* 2010; **54**: 147–53.
- Hu B, Bao R, Wang X, et al. The size of endotracheal tube and sore throat after surgery: a systematic review and meta-analysis. *PLoS ONE* 2013; **8**: e74467.
- Koh KF, Hare JD, Calder I. Small tubes revisited. *Anaesthesia* 1998; **53**: 46–50.
- Najafi A, Imani F, Makarem J, et al. Postoperative sore throat after laryngoscopy with macintosh or glide scope video laryngoscope blade in normal airway patients. *Anesthesiology and Pain Medicine* 2014; **4**: e15136.
- Maruyama K, Nakagawa H, Imanishi H, Kitamura A, Haya-shida M. Comparison of postoperative pharyngeal morbidity using the Macintosh laryngoscope or AirWay Scope after mastectomy. *Journal of Anesthesia* 2011; **25**: 773.
- Combes X, Andriamifidy L, Dufresne E, et al. Comparison of two induction regimens using or not using muscle relaxant:

- impact on postoperative upper airway discomfort. *British Journal of Anaesthesia* 2007; **99**: 276–81.
12. Mencke T, Knoll H, Schreiber JU, et al. Rocuronium is not associated with more vocal cord injuries than succinylcholine after rapid-sequence induction: a randomized, prospective, controlled trial. *Anesthesia and Analgesia* 2006; **102**: 943–9.
 13. Clayton-Smith A, Bennett K, Alston RP, et al. A comparison of the efficacy and adverse effects of double-lumen endobronchial tubes and bronchial blockers in thoracic surgery: a systematic review and meta-analysis of randomized controlled trials. *Journal of Cardiothoracic and Vascular Anesthesia* 2015; **29**: 955–66.
 14. Jeon J, Lee K, Ahn G, Lee J, Hwang W. Comparison of postoperative sore throat and hoarseness between two types of double-lumen endobronchial tubes: a randomized controlled trial. *Journal of Cardiothoracic and Vascular Anesthesia* 2015; **29**: 121–5.
 15. Seo JH, Kwon TK, Jeon Y, Hong DM, Kim HJ, Bahk JH. Comparison of techniques for double-lumen endobronchial intubation: 90° or 180° rotation during advancement through the glottis. *British Journal of Anaesthesia* 2013; **111**: 812.
 16. Zhong T, Wang W, Chen J, Ran L, Story DA. Sore throat or hoarse voice with bronchial blockers or double-lumen tubes for lung isolation: a randomised, prospective trial. *Anaesthesia and Intensive Care* 2009; **37**: 441.
 17. Turkstra TP, Smitheram AK, Alabdulhadi O, Youssef H, Jones PM. The Flex-Tip™ tracheal tube does not reduce the incidence of postoperative sore throat: a randomized controlled trial. *Canadian Journal of Anesthesia* 2011; **58**: 1090–6.
 18. Tanaka Y, Nakayama T, Nishimori M, Tsujimura Y, Kawaguchi M, Sato Y. Lidocaine for preventing postoperative sore throat. *Cochrane Database of Systematic Reviews* 2015; **7**: Cd004081.
 19. Banihashem N, Alijanpour E, Hasannasab B, Zarei A. Prophylactic effects of lidocaine or beclomethasone spray on postoperative sore throat and cough after orotracheal intubation. *Iranian Journal of Otorhinolaryngology* 2015; **27**: 179–84.
 20. Bousselmi R, Lebbi MA, Bargaoui A, et al. Lidocaine reduces endotracheal tube associated side effects when instilled over the glottis but not when used to inflate the cuff: a double blind, placebo-controlled, randomized trial. *Tunisie Medicale* 2014; **92**: 29–33.
 21. Suma KV, Udaya K. Prevention of post intubation sore throat by inflating endotracheal tube cuff with alkalinized lignocaine. *Indian Journal of Public Health Research and Development* 2015; **6**: 200–4.
 22. Sun L, Guo R, Sun L. Dexamethasone for preventing postoperative sore throat: a meta-analysis of randomized controlled trials. *Irish Journal of Medical Science* 2014; **183**: 593–600.
 23. Park S, Kim S, Lee S, et al. Application of triamcinolone acetonide paste to the endotracheal tube reduces postoperative sore throat: a randomized controlled trial. *Canadian Journal of Anesthesia* 2011; **58**: 436–42.
 24. Sumathi PA, Shenoy T, Ambaresha M, Krishna HM. Controlled comparison between betamethasone gel and lidocaine jelly applied over tracheal tube to reduce postoperative sore throat, cough, and hoarseness of voice. *British Journal of Anaesthesia* 2008; **100**: 215–8.
 25. Tazeh-kand NF, Eslami B, Mohammadian K. Inhaled fluticasone propionate reduces postoperative sore throat, cough, and hoarseness. *Anesthesia and Analgesia* 2010; **111**: 895–8.
 26. Chen Y-Q, Li J-P, Xiao J. Prophylactic effectiveness of budesonide inhalation in reducing postoperative throat complaints. *European Archives of Oto-rhino-laryngology* 2014; **271**: 1667.
 27. Chen C-Y, Kuo C-J, Lee Y-W, Lam F, Tam K-W. Benzylamine hydrochloride on postoperative sore throat: a meta-analysis of randomized controlled trials. *Canadian Journal of Anesthesia* 2014; **61**: 220–8.
 28. Chang JE, Min SW, Kim CS, Han SH, Kwon YS, Hwang JY. Effect of prophylactic benzylamine hydrochloride on postoperative sore throat and hoarseness after tracheal intubation using a double-lumen endobronchial tube: a randomized controlled trial. *Canadian Journal of Anesthesia* 2015; **62**: 1097–103.
 29. Agarwal A, Nath SS, Goswami D, Gupta D, Dhiraaj S, Singh PK. An evaluation of the efficacy of aspirin and benzylamine hydrochloride gargle for attenuating postoperative sore throat: a prospective, randomized. *Anesthesia and Analgesia* 2006; **103**: 1001–3.
 30. Thang'a P, Kamya D, Mung'ayi V. Effects of intravenous diclofenac on postoperative sore throat in patients undergoing laparoscopic surgery at Aga Khan University Hospital, Nairobi: a prospective, randomized, double blind controlled trial. *African Health Sciences* 2013; **13**: 999.
 31. Rahimi M, Makarem J. Effects of diclofenac epolamine patch on postoperative sore throat in parturients after cesarean delivery under endotracheal general anesthesia. *Acta Anaesthesiologica Taiwanica* 2009; **47**: 17.
 32. Liu J, Zhang X, Gong W, et al. Correlations between controlled endotracheal tube cuff pressure and postprocedural complications: a multicenter study. *Anesthesia and Analgesia* 2010; **111**: 1133–7.
 33. Ansari L, Bohluli B, Mahaseni H, Valaei N, Sadr-Eshkevari P, Rashad A. The effect of endotracheal tube cuff pressure control on postextubation throat pain in orthognathic surgeries: a randomized double-blind controlled clinical trial. *British Journal of Oral and Maxillofacial Surgery* 2014; **52**: 140–3.
 34. Ryu J-H, Han S-S, Do S-H, Lee J-M, Lee S-C, Choi E-S. Effect of adjusted cuff pressure of endotracheal tube during thyroidectomy on postoperative airway complications: prospective, randomized, and controlled trial. *World Journal of Surgery* 2013; **37**: 786.
 35. Agarwal A, Gupta D, Yadav G, Goyal P, Singh PK, Singh U. An evaluation of the efficacy of licorice gargle for attenuating postoperative sore throat: a prospective, randomized, single-blind study. *Anesthesia and Analgesia* 2009; **109**: 77–81.
 36. Ruetzler K, Fleck M, Nabecker S, et al. A randomized, double-blind comparison of licorice versus sugar-water gargle for prevention of postoperative sore throat and postextubation coughing. *Anesthesia and Analgesia* 2013; **117**: 614–21.
 37. Gupta D, Agrawal S, Sharma JP. Effect of preoperative licorice lozenges on incidence of postextubation cough and sore throat in smokers undergoing general anesthesia and endotracheal intubation. *Middle East Journal of Anaesthesiology* 2013; **22**: 173–8.
 38. Borazan H, Kecicioglu A, Okesli S, Otelcioglu S. Oral magnesium lozenge reduces postoperative sore throat: a randomized, prospective, placebo-controlled study. *Anesthesiology* 2012; **117**: 512–8.
 39. Teymourian H, Mohajerani SA, Farahbod A. Magnesium and ketamine gargle and postoperative sore throat. *Anesthesiology and Pain Medicine* 2015; **5**: e22367.
 40. Canbay O, Celebi N, Sahin A, Celiker V, Ozgen S, Aypar U. Ketamine gargle for attenuating postoperative sore throat. *British Journal of Anaesthesia* 2008; **100**: 490–3.

41. Park SY, Kim SH, Noh JJ, et al. The effect of intravenous low dose ketamine for reducing postoperative sore throat. *Korean Journal of Anesthesiology* 2010; **59**: 22–6.
42. Ebneshahidi A, Mohseni M. Strepsils® tablets reduce sore throat and hoarseness after tracheal intubation. *Anesthesia and Analgesia* 2010; **111**: 892.
43. Gulhas N, Canpolat H, Cicek M, et al. Dexpanthenol pastille and benzydamine hydrochloride spray for the prevention of post-operative sore throat. *Acta Anaesthesiologica Scandinavica* 2007; **51**: 239.
44. Gupta D, Agrawal S, Sharma JP. Evaluation of preoperative Strepsils lozenges on incidence of postextubation cough and sore throat in smokers undergoing anesthesia with endotracheal intubation. *Saudi Journal of Anaesthesia* 2014; **8**: 244–8.
45. Ogata J, Minami K, Horishita T, et al. Gargling with sodium azulene sulfonate reduces the postoperative sore throat after intubation of the trachea. *Anesthesia and Analgesia* 2005; **101**: 290.
46. Maruyama K, Yamada T, Hara K. Effect of clonidine premedication on postoperative sore throat and hoarseness after total intravenous anesthesia. *Journal of Anesthesia* 2006; **20**: 327–30.
47. Cook T, Howes B. Supraglottic airway devices: recent advances. *Continuing Education in Anaesthesia, Critical Care and Pain* 2011; **11**: 56–61.
48. Cook T, Woodall N, Frerk C. 4th National Audit Project of the Royal College of Anaesthetists and The Difficult Airway Society. *Anesthesiology* 2011; **116**: 216.
49. Wong DT, Tam AD, Mehta V, Raveendran R, Riad W, Chung FF. New supraglottic airway with built-in pressure indicator decreases postoperative pharyngolaryngeal symptoms: a randomized controlled trial. *Canadian Journal of Anesthesia* 2013; **60**: 1197–203.
50. Keijzer C, Buitelaar DR, Efthymiou KM, et al. A comparison of postoperative throat and neck complaints after the use of the i-gel and the La Premiere disposable laryngeal mask: a double-blinded, randomized, controlled trial. *Anesthesia and Analgesia* 2009; **109**: 1092–5.
51. Uerpairojkit K, Charuluxananan S, Werawatganon T, Poomseetong T. Profile Soft-Seal Cuff for general anesthesia under ambulatory gynecologic laparoscopy. *Journal of the Medical Association of Thailand* 2009; **92**: 1184–90.
52. Ng SY, Teoh WHL, Lim Y, Cheong VG. Comparison of the AMBU laryngeal mask and the LMA laryngeal mask classic in anaesthetised, spontaneously breathing patients. *Anaesthesia and Intensive Care* 2007; **35**: 57–61.
53. Brimacombe J, Holyoake L, Keller C, et al. Emergence characteristics and postoperative laryngopharyngeal morbidity with the laryngeal mask airway: a comparison of high versus low initial cuff volume. *Anaesthesia* 2000; **55**: 338–43.
54. Tan MGE, Chin ERC, Kong CS, Chan YH, Ip-Yam PC. Comparison of the re-usable LMA laryngeal mask Classic and two single-use laryngeal masks (LMA laryngeal mask Unique and SoftSeal) in airway management by novice personnel. *Anaesthesia and Intensive Care* 2005; **33**: 739–43.
55. van Zundert AAJ, Fonck K, Al-Shaikh B, Mortier EP. Comparison of cuff-pressure changes in LMA laryngeal mask Classic and the new Soft Seal laryngeal masks during nitrous oxide anaesthesia in spontaneous breathing patients. *European Journal of Anaesthesiology* 2004; **21**: 547–52.
56. Van Zundert AAJ, Fonck K, Al-Shaikh B, Mortier E. Comparison of the LMA laryngeal mask-classic with the new disposable soft seal laryngeal mask in spontaneously breathing adult patients. *Anesthesiology* 2003; **99**: 1066–71.
57. Paech MJ, Lain J, Garrett WR, Gillespie G, Stannard KJD, Doherty DA. Randomized evaluation of the single-use SoftSeal and the re-useable LMA laryngeal mask Classic laryngeal mask. *Anaesthesia and Intensive Care* 2004; **32**: 66–72.
58. Cook TM, Trümpelmann P, Beringer R, Stedeford J. A randomized comparison of the Portex Softseal™ laryngeal mask airway with the LMA laryngeal mask-Unique™ during anaesthesia. *Anaesthesia* 2005; **60**: 1218–25.
59. van Zundert AAJ, Al-Shaikh B, Brimacombe J, Koster J, Koning D, Mortier E. Comparison of three disposable extraglottic airway devices in spontaneously breathing adults. *Anesthesiology* 2006; **104**: 1165–9.
60. Turan A, Kaya G, Koyuncu O, Karamanlioglu B, Pamukçu Z. Comparison of the laryngeal mask (LMA laryngeal mask) and laryngeal tube (LT) with the new perilaryngeal airway (CobraPLA) in short surgical procedures. *European Journal of Anaesthesiology* 2006; **23**: 234–8.
61. Gaitini L, Yanovski B, Somri M, Vaida S, Riad T, Alfery D. A comparison between the PLA Cobra and the Laryngeal Mask Airway Unique during spontaneous ventilation: a randomized prospective study. *Anesthesia and Analgesia* 2006; **102**: 631–6.
62. Yaghoobi S, Abootorabi SM, Kayalha H, Van Zundert TC, Pakpour AH. Efficacy of the new perilaryngeal airway (Cobra-PLA™) versus the laryngeal mask airway (LMA™) to improve oropharyngeal leak pressure in obese and overweight patients. *Tanaffos* 2015; **14**: 42–8.
63. de Montblanc J, Ruscio L, Maziot JX, Benhamou D. A systematic review and meta-analysis of the i-gel® vs laryngeal mask airway in adults. *Anaesthesia* 2014; **69**: 1151–62.
64. Park SK, Choi GJ, Choi YS, Ahn EJ, Kang H. Comparison of the i-gel and the laryngeal mask airway proSeal during general anesthesia: a systematic review and meta-analysis. *PLoS ONE* 2015; **10**: e0119469.
65. Chen X, Jiao J, Cong X, Liu L, Wu X. A comparison of the performance of the i-gel vs. the LMA laryngeal mask-S during anesthesia: a meta-analysis of randomized controlled trials. *PLoS ONE* 2013; **8**: e71910.
66. Francksen H, Renner J, Hanss R, Scholz J, Doerges V, Bein B. A comparison of the i-gel with the LMA laryngeal mask-Unique in non-paralysed anaesthetised adult patients. *Anaesthesia* 2009; **64**: 1118–24.
67. Maitra S, Khanna P, Baidya DK. Comparison of laryngeal mask airway Supreme and laryngeal mask airway Pro-Seal for controlled ventilation during general anaesthesia in adult patients: systematic review with meta-analysis. *European Journal of Anaesthesiology* 2014; **31**: 266–73.
68. Pajiyar AK, Wen Z, Wang H, Ma L, Miao L, Wang G. Comparisons of clinical performance of guardian laryngeal mask with laryngeal mask airway ProSeal. *BMC Anesthesiology* 2015; **15**: 69.
69. Shin W-J, Cheong Y-S, Yang H-S, Nishiyama T. The supraglottic airway I-gel in comparison with ProSeal laryngeal mask airway and classic laryngeal mask airway in anaesthetized patients. *European Journal of Anaesthesiology* 2010; **27**: 598–601.
70. Jadhav PA, Dalvi NP, Tendolkar BA. I-gel versus laryngeal mask airway-ProSeal: comparison of two supraglottic airway devices in short surgical procedures. *Journal of Anaesthesiology, Clinical Pharmacology* 2015; **31**: 221–5.
71. Miller DM, Camporota L. Advantages of ProSeal and SLIPA airways over tracheal tubes for gynecological laparoscopies. *Canadian Journal of Anesthesia* 2006; **53**: 188–93.

72. Choi YM, Cha SM, Kang H, et al. The clinical effectiveness of the streamlined liner of pharyngeal airway (SLIPA) compared with the laryngeal mask airway ProSeal during general anaesthesia. *Korean Journal of Anesthesiology* 2010; **58**: 450–7.
73. Ali A, Canturk S, Turkmen A, Turgut N, Altan A. Comparison of the laryngeal mask airway Supreme and laryngeal mask airway Classic in adults. *European Journal of Anaesthesiology* 2009; **26**: 1010–4.
74. Lange M, Smul T, Zimmermann P, Kohlenberger R, Roewer N, Kehl F. The effectiveness and patient comfort of the novel streamlined pharynx airway liner (SLIPA) compared with the conventional laryngeal mask airway in ophthalmic surgery. *Anesthesia and Analgesia* 2007; **104**: 431–4.
75. Baidya DK, Chandrakha Darlong V, Pandey R, Maitra S, Khanna P. Comparative efficacy and safety of the Ambu® AuraOnce™ laryngeal mask airway during general anaesthesia in adults: a systematic review and meta-analysis. *Anaesthesia* 2014; **69**: 1023–32.
76. Donaldson W, Abraham A, Deighan M, Michalek P. i-gel™ vs. AuraOnce™ laryngeal mask for general anaesthesia with controlled ventilation in paralyzed patients. *Biomedical papers of the Medical Faculty of the University Palacký, Olomouc, Czechoslovakia* 2011; **55**: 155–63.
77. Middleton P. Insertion techniques of the laryngeal mask airway: a literature review. *Journal of Perioperative Practice* 2009; **19**: 31–5.
78. Krishna HM, Kamath S, Shenoy L. Insertion of LMA laryngeal mask Classic™ with and without digital intraoral manipulation in anesthetized unparalyzed patients. *Journal of Anaesthesiology and Clinical Pharmacology* 2012; **28**: 481–5.
79. Bergmann I, Crozier TA, Roessler M, et al. The effect of changing the sequence of cuff inflation and device fixation with the LMA laryngeal mask-Supreme® on device position, ventilatory complications, and airway morbidity: a clinical and fiberoptic study. *BMC Anesthesiology* 2014; **14**: 2.
80. Choo CY, Koay CK, Yoong CS. A randomised controlled trial comparing two insertion techniques for the Laryngeal Mask Airway Flexible in patients undergoing dental surgery. *Anaesthesia* 2012; **67**: 986–90.
81. Hwang J, Park H, Lim Y, Do S, Lee S, Jeon Y. Comparison of two insertion techniques of ProSeal™ Laryngeal Mask Airway. *Anesthesiology* 2009; **110**: 905–7.
82. Jeon YT, Na HS, Park SH, et al. Insertion of the ProSeal laryngeal mask airway is more successful with the 90 degree rotation technique. *Canadian Journal of Anesthesia* 2010; **57**: 211–5.
83. Kuppasamy A, Azhar N. Comparison of bougie-guided insertion of ProSeal™ laryngeal mask airway with digital technique in adults. *Indian Journal of Anaesthesia* 2010; **54**: 35–9.
84. Maclean J, Tripathy D, Parthasarathy S, Ravishankar M. Comparative evaluation of gum-elastic bougie and introducer tool as aids in positioning of ProSeal laryngeal mask airway in patients with simulated restricted neck mobility. *Indian Journal of Anaesthesia* 2013; **57**: 248–52.
85. Chen HS, Yang SC, Chien CFC, Spielberger J, Hung KC, Chung KC. Insertion of the ProSeal™ laryngeal mask airway is more successful with the Flexi-Slip™ stylet than with the introducer. *Canadian Journal of Anesthesia* 2011; **58**: 617–23.
86. Chen M, Hsu H, Lu I, Shih C, Shen Y, Tseng K. Techniques for the insertion of the ProSeal laryngeal mask airway: comparison of the Foley airway stylet tool with the introducer tool in a prospective, randomized study. *BMC Anesthesiology* 2014; **14**: 1–7.
87. García-Aguado R, Viñoles J, Brimacombe J, Vivó M, López-Estudillo R, Ayala G. suction catheter guided insertion of the ProSeal laryngeal mask airway is superior to the digital technique. *Canadian Journal of Anesthesia* 2006; **53**: 398–403.
88. Chantzara G, Stroumpoulis K, Alexandrou N, Kokkinos L, Iacovidou N, Xanthos T. Influence of LMA laryngeal mask cuff pressure on the incidence of pharyngolaryngeal adverse effects and evaluation of the use of manometry during different ventilation modes: a randomized clinical trial. *Minerva Anestesiologica* 2014; **80**: 547–55.
89. Seet E, Yousaf F, Gupta S, Subramanyam R, Wong DT, Chung F. Use of manometry for laryngeal mask airway reduces postoperative pharyngolaryngeal adverse events: a prospective, randomized trial. *Anesthesiology* 2010; **112**: 652–7.
90. Jeon Y-S, Choi J-W, Jung H-S, et al. Effect of continuous cuff pressure regulator in general anaesthesia with laryngeal mask airway. *Journal of International Medical Research* 2011; **39**: 1900–7.
91. Dostalova K, Doubravska L, Fritscherova S, et al. Monitoring sore throat after the introduction of laryngeal masks. *Anesteziologie a Intenzivni Medicina* 2015; **26**: 72–8.
92. Gunenc F, Kucukguclu S, Ozbilgin S, Kuvaki B, Maltepe F. Effect of laryngeal mask cuff pressure on postoperative pharyngolaryngeal morbidity in geriatric patients. *Turk Geriatri Dergisi* 2015; **18**: 143–50.
93. Li BB, Yan J, Zhou HG, Hao J, Liu AJ, Ma ZL. Application of minimum effective cuff inflating volume for laryngeal mask airway and its impact on postoperative pharyngeal complications. *Chinese Medical Journal* 2015; **128**: 2570–6.
94. Vasanth Karthik R, Ranganathan P, Kulkarni AP, Sharma KS. Does cuff pressure monitoring reduce postoperative pharyngolaryngeal adverse events after LMA laryngeal mask-ProSeal insertion? A parallel group randomised trial. *Journal of Anesthesia* 2014; **28**: 662–7.
95. Joe HB, Kim DH, Chae YJ, Kim JY, Kang M, Park KS. The effect of cuff pressure on postoperative sore throat after Cobra perilaryngeal airway. *Journal of Anesthesia* 2012; **26**: 225–9.
96. Kang JE, Oh CS, Choi JW, Son IS, Kim SH. Postoperative pharyngolaryngeal adverse events with laryngeal mask airway (LMA laryngeal mask Supreme) in laparoscopic surgical procedures with cuff pressure limiting 25 cm H₂O: prospective, blind, and randomised study. *Scientific World Journal* 2014; **2014**: 709–801.
97. Nasserri K, Shami S, Rashidzadeh L, Kanisiran LA, Care I, Practitioner G. Evaluation of lidocaine efficacy on attenuation of sore throat, cough and itching after laryngeal mask airway insertion in elective eye surgery. *Journal of Mazandaran University of Medical Sciences* 2013; **23**: 103–7.
98. Kiran S, Goel M, Singhal P, Gupta N, Bhardwaj M. Postoperative sore throat with 0.05% betamethasone gel and 2% lignocaine jelly used as a lubricant for ProSeal LMA laryngeal mask (PLMA laryngeal mask) insertion. *Egyptian Journal of Anaesthesia* 2012; **28**: 139–42.
99. Taghavi Gilani M, Miri Soleimani I, Razavi M, Salehi M. Reducing sore throat following laryngeal mask airway insertion: comparing lidocaine gel, saline, and washing mouth with the control group. *Revista Brasileira de Anestesiologia* 2013; **65**: 450–4.
100. Kim DK, Rhee KY, Kwon WK, Kim TY, Kang JE. A heated humidifier does not reduce laryngo-pharyngeal complaints after brief laryngeal mask anaesthesia. *Canadian Journal of Anesthesia* 2007; **54**: 134–40.

101. Uztire N, Menda F, Bilgen S, Keskin O, Temur S, Koner O. The effect of flurbiprofen on postoperative sore throat and hoarseness after LMA laryngeal mask-ProSeal insertion: a randomised, clinical trial. *Turkish Journal of Anesthesia and Reanimation* 2014; **42**: 123–7.
102. Rashwan S, Abdelmawgoud A, Badawy A. Effect of tramadol gargle on postoperative sore throat: a double blinded randomized placebo controlled study. *Egyptian Journal of Anaesthesia* 2014; **30**: 235–9.
103. Eydi M, Kolahdouzan K, Golzari SE. Effect of intravenous hydrocortisone on preventing postoperative sore throat followed by laryngeal mask airway use in patients undergoing urogenital surgeries. *Journal of Cardiovascular and Thoracic Research* 2013; **5**: 29–33.
104. Maino P, Dullenkopf A, Bernet V, Weiss M. Nitrous oxide diffusion into the cuffs of disposable laryngeal mask airways. *Anaesthesia* 2005; **60**: 278–82.
105. Na H-S, Jeon Y-T, Shin H-J, Oh A-Y, Park H-P, Hwang J-W. Effect of paralysis at the time of proseal laryngeal mask airway insertion on pharyngolaryngeal morbidities. A randomized trial. *PLoS ONE* 2015; **10**: e0134130.
106. Chia YY, Lee SW, Liu K. Propofol causes less postoperative pharyngeal morbidity than thiopental after the use of a laryngeal mask airway. *Anesthesia and Analgesia* 2008; **106**: 123–6.
107. Metry AA. Does the anesthetic agent used for maintenance of anesthesia affect post-operative pharyngeal and laryngeal morbidities after laryngeal mask airway insertion? *Acta Anaesthesiologica Italica* 2009; **60**: 45–56.
108. The Royal College of Anaesthetists. Your child's general anaesthetic. http://www.rcoa.ac.uk/system/files/PI-YCGA-COL-2014_0.pdf (accessed 15/01/2016).
109. The Evelina London Children's Hospital. Your child's general anaesthetic, 2013. <http://www.evelinalondon.nhs.uk/resources/patient-information/your-childs-anaesthetic.pdf> (accessed 15/01/2016).
110. Wong DDL, Bailey CR. Emergence delirium in children. *Anaesthesia* 2015; **70**: 36–8.
111. Tekelioglu UY, Apuhan T, Akkaya A, et al. Comparison of topical tramadol and ketamine in pain treatment after tonsillectomy. *Pediatric Anesthesia* 2013; **23**: 496–501.
112. Harding CJ, McVey FK. Interview method affects incidence of postoperative sore throat. *Anaesthesia* 1987; **42**: 1104–7.
113. Stout DM, Bishop MJ, Dwersteg JF, Cullen BF. Correlation of endotracheal tube size with sore throat and hoarseness following general anesthesia. *Anesthesiology* 1987; **67**: 419–21.
114. Beringer RM, Kelly F, Cook TM, et al. A cohort evaluation of the paediatric i-gel™ airway during anaesthesia in 120 children. *Anaesthesia* 2011; **66**: 1121–6.
115. von Ungern-Sternberg BS, Ramgolam A, Hall GL, Sly PD, Habre W. Peri-operative adverse respiratory events in children. *Anaesthesia* 2015; **70**: 440–4.
116. Huang HJ, Fang XM. Effect of endotracheal intubation and laryngeal mask airway on perioperative respiratory adverse events in children with upper airway infections. *Zhonghua Yi Xue Za Zhi* 2013; **93**: 3626–8.
117. Gharai B, Aghamohammadi H, Jafari A, Razavi S, Kamranmanesh M, Kermany AS. Use of laryngeal mask airway in children with upper respiratory tract infection, compared with face mask: randomized, single blind, clinical trial. *Acta Anaesthesiologica Taiwan* 2011; **49**: 136–40.
118. Patel MG, Swadia V, Bansal G. Prospective randomized comparative study of use of PLMA laryngeal mask and ET tube for airway management in children under general anaesthesia. *Indian Journal of Anaesthesia* 2010; **54**: 109–15.
119. Hara K, Maruyama K. Effect of additives in lidocaine spray on postoperative sore throat, hoarseness and dysphagia after total intravenous anaesthesia. *Acta Anaesthesiologica Scandinavica* 2005; **49**: 463–7.
120. Borazan H, Sarıtaş TB. Predictors of postoperative sore throat in intubated children. *Pediatric Anesthesia* 2012; **22**: 595–6.
121. Zhao N, Deng F, Yu C. Anesthesia for pediatric day-case dental surgery: a study comparing the classic laryngeal mask airway with nasal tracheal intubation. *Journal of Craniofacial Surgery* 2014; **25**: e245–8.
122. Patki A. Laryngeal mask airway vs the endotracheal tube in paediatric airway management: a meta-analysis of prospective randomised controlled trials. *Indian Journal of Anaesthesia* 2011; **55**: 537–41.
123. Luce V, Harkouk H, Brasher C, et al. Supraglottic airway devices vs tracheal intubation in children: a quantitative meta-analysis of respiratory complications. *Pediatric Anesthesia* 2014; **24**: 1088–98.
124. Needleman HL, Harpavat S, Wu S, Allred EN, Berde C. Postoperative pain and other sequelae of dental rehabilitations performed on children under general anesthesia. *Pediatric Dentistry* 2008; **30**: 111–21.
125. Calder A, Hegarty M, Erb TO, von Ungern-Sternberg BS. Predictors of postoperative sore throat in intubated children. *Pediatric Anesthesia* 2012; **22**: 239–43.
126. Das B, Mitra S, Jamil SN, Varshney RK. Comparison of three supraglottic devices in anesthetized paralyzed children undergoing elective surgery. *Saudi Journal of Anaesthesia* 2012; **6**: 224–8.
127. Kim H, Lee JY, Lee SY, Park SY, Lee SC, Chung CJ. A comparison of i-gel™ and LMA laryngeal mask Supreme™ in anesthetized and paralyzed children. *Korean Journal of Anesthesiology* 2014; **67**: 317–22.
128. Drake-Brockman TFE, Ledowski T, Hegarty M, Gessner M, von Ungern-Sternberg BS. A comparison of the i-gel™ and the PRO-Breathe® Laryngeal Mask during pressure support ventilation in children. *Anaesthesia* 2015; **70**: 1412–7.
129. Smith P, Bailey CR. A performance comparison of the paediatric i-gel® with other supraglottic airway devices. *Anaesthesia* 2015; **70**: 84–92.
130. Hockings L, Heaney M, Chambers NA, Erb TO, von Ungern-Sternberg BS. Reduced air leakage by adjusting the cuff pressure in pediatric laryngeal mask airways during spontaneous ventilation. *Pediatric Anesthesia* 2010; **20**: 313–7.
131. Wong JG, Heaney M, Chambers NA, Erb TO, von Ungern-Sternberg BS. Impact of laryngeal mask airway cuff pressures on the incidence of sore throat in children. *Pediatric Anesthesia* 2009; **19**: 464–9.
132. Von Ungern-Sternberg BS, Erb TO, Chambers NA, Heaney M. Laryngeal mask airways - - to inflate or to deflate after insertion? *Pediatric Anesthesia* 2009; **19**: 837–43.
133. Chen BZ, Luo LH, Jiang L, Wang RR, Li J, Tan L. The effect of nitrous oxide on intracuff pressure of the size 2 ProSeal laryngeal Mask Airway. *Journal of Clinical Anesthesia* 2011; **23**: 214–7.
134. William A, Chambers NA, Erb TO, von-Ungern-Sternberg BS. Incidence of sore throat in children following use of flexible laryngeal mask airways – impact of an introducer device. *Pediatric Anesthesia* 2010; **20**: 839–43.